

Impact of continuous NOMAC/E₂ (Zoely®) regimen on quality of life in women with dysmenorrhea and sonographic suspicion of endometriosis

Josep Perelló-Capó¹, Bruna Miarons¹, Natalia Teixeira¹, Noelia Rams¹, Rocío Luna¹, Josep Estadella¹, Taisiia Melnychuk¹, Ramon Rovira¹

¹ Department of Obstetrics and Gynecology, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

ABSTRACT

Background: Women with endometriosis have a lower quality of life (QoL) than those without the disease. Treatment should therefore aim not only to improve symptoms but also to enhance overall QoL.

Objective: To evaluate QoL in patients with dysmenorrhea and ultrasound suspicion of endometriosis after starting nomegestrol acetate 2.5 mg plus 17 β -estradiol 1.5 mg (NOMAC/E₂) in an extended regimen.

Methods: Prospective cohort pilot study evaluating changes in QoL in women with ultrasound-suspected endometriosis treated with NOMAC/E₂ for dysmenorrhea. QoL was assessed using the EuroQoL-5D questionnaire, including the EQ-5D index and the visual numeric scale (EQ-VAS). Changes after initiating treatment were analyzed using a repeated-measures general linear model and confirmed with the Wilcoxon test.

Results: Fourteen patients were included (mean \pm SD age 36.6 \pm 7.4 years). Significant improvements were observed in QoL after starting treatment. The EQ-5D increased from 0.65 \pm 0.22 at baseline to 0.86 \pm 0.11 at 6 months and 0.93 \pm 0.07 at 12 months ($p < 0.001$). The EQ-VAS index increased from 78.4 \pm 12.7 points at baseline to 89.2 \pm 9.7 points at 6 months and 94.4 \pm 4.0 points at 12 months ($p < 0.001$).

Conclusions: Continuous treatment with NOMAC/E₂ in women with dysmenorrhea and ultrasound suspicion of endometriosis may significantly improve QoL as measured with the EQ-5D questionnaire.

KEYWORDS

Endometriosis, quality of life, EQ-5D, dysmenorrhea, nomegestrol acetate, 17 β -estradiol, NOMAC/E₂, combined hormonal contraceptive.

Introduction

Endometriosis is a condition in which endometrial glands and stroma develop outside the uterine cavity^[1]. It is common, affecting 5–10% of women of reproductive age^[2–4], and may appear as peritoneal lesions, endometriomas, or deep infiltrating endometriosis (DIE). Typical symptoms include dysmenorrhea, dyschezia, dyspareunia, dysuria, chronic pelvic pain, and infertility. Beyond physical discomfort, endometriosis can significantly affect psychological health, social functioning, sexual satisfaction, and overall quality of life (QoL)^[5–10].

Management includes both pharmacological and surgical options. Treatment should be individualized according to symptom profile, severity, reproductive desire, age, extent and location of disease, and response to previous therapies^[11]. Surgery was historically considered the primary treatment. However, concerns over surgical risk, the potential negative impact on ovarian reserve^[12–14], the high recurrence rate after surgery^[15], and the expanding availability of effective hormonal therapies have shifted practice. For women not seeking conception who present with clinical and imaging findings suggestive of endometriosis, hormonal treatment—particularly combined hormonal contraceptives—has become the preferred first-line strategy for pain control^[13]. Despite this shift, evidence guiding the optimal choice of product and regimen for specific lesion types and patient characteristics remains limited.

Article history

Received 30 Jun 2025 - Accepted 20 Aug 2025

Contact

Josep Perelló Capó; JPerello@santpau.cat
Attending Physician Obstetrics and Gynecology
Hospital de la Santa Creu i Sant Pau
Autonomous University of Barcelona
Barcelona, Spain

DOI

10.53260/E60.257034

Treatment response varies widely, from complete symptom resolution to minimal improvement, and symptoms typically recur if therapy is discontinued^[16]. Consequently, long-term hormonal treatment is recommended, ideally continuing until pregnancy is desired or menopause occurs^[17]. In addition to symptom relief and prevention of recurrence, improving QoL is a fundamental goal.

Nomegestrol acetate 2.5 mg combined with 17 β -estradiol 1.5 mg (NOMAC/E₂)—a formulation pairing a strongly antiproliferative progestogen with a natural estrogen of low proliferative activity—has emerged as a promising option. The Spanish Fertility Society (Sociedad Española de Fertilidad) includes NOMAC/E₂ among recommended treatments for patients with endometriosis^[18].

The primary objective of this study was to assess whether patients with dysmenorrhea and ultrasound findings suggestive of endometriosis (endometrioma ± deep infiltrating endometriosis [DIE]) experienced improvement in QoL after initiating NOMAC/E₂. Secondary objectives included evaluating changes in the intensity and frequency of dysmenorrhea, non-cyclic chronic pelvic pain (CPP), dyschezia, dysuria, and dyspareunia; changes in menstrual bleeding patterns; ultrasound-measured changes in lesion size and DIE; patient satisfaction; and treatment-related side effects.

Methods

This was a prospective cohort pilot study including all patients referred from primary care to our tertiary hospital (Hospital de la Santa Creu i Pau, Barcelona, Spain) with ultrasound findings suggestive of endometriosis who received NOMAC/E₂ between 01/01/2016 and 31/03/2019 for dysmenorrhea management. The study was approved by the Ethics Committee of Hospital de la Santa Creu i Sant Pau (Barcelona, Spain).

Eligible participants were women of reproductive age with regular menstrual cycles (24–38 days), dysmenorrhea ≥ 3 on a visual numeric scale (VNS), and ultrasound findings consistent with endometriosis based on IDEA criteria. The IDEA protocol includes standardized transvaginal assessment of the uterus and adnexa (for adenomyosis and endometriomas), evaluation of ovarian mobility and site-specific tenderness, inspection of the pouch of Douglas for obliteration, and systematic identification of deep endometriosis nodules in the anterior and posterior compartments. Additional inclusion criteria were absence of pregnancy, no hormonal therapy in the preceding 3 months, and no intention to conceive within the next year. Patients with contraindications to combined hormonal contraceptives (WHO eligibility criteria) were excluded. All participants provided written informed consent. Although NOMAC/E₂ is typically administered in a 24/4 regimen, all patients in this study were instructed to use continuous dosing without hormone-free intervals.

During the study period, 187 women were evaluated for suspected endometriosis; 14 met inclusion criteria and started continuous NOMAC/E₂ (Zoely®).

The primary outcome was QoL, assessed using the EuroQol-5D (EQ-5D) questionnaire^[19]. The EQ-5D includes a descriptive system covering five domains (mobility, self-care, usual activities, pain/discomfort, and anxiety/depression), each with five severity levels, yielding a global EQ-5D health state score. Patients also rated their overall health using the EQ visual analogue scale (EQ-VAS), a 20-cm vertical scale from 0 (worst imaginable health) to 100 (best imaginable health). EQ-5D scores were calculated using reference values established for our population^[20].

Secondary outcomes included duration and intensity of dysmenorrhea, menstrual bleeding patterns, chronic pelvic pain (CPP), dysuria, dyschezia, and dyspareunia. Pain duration was measured in days, and intensity was rated using a VNS ranging from

0 (no pain) to 10 (worst pain imaginable). Patients reported the average number of days with pain, pain intensity, and bleeding days based on the previous three menstrual cycles. Menstrual bleeding intensity was classified using a 4-item Likert scale (amenorrhea, light, normal, abundant).

Ultrasound outcomes included maximum diameter of the largest endometrioma, maximum diameter of the largest DIE lesion, and the sum of maximum diameters of all endometriomas. Need for surgery, treatment changes, adverse effects, and satisfaction were also recorded. Treatment satisfaction was assessed using a VNS from 0 (completely dissatisfied) to 5 (completely satisfied). Participants attended three visits: baseline (V0), 6-month follow-up (V1), and 12-month follow-up (V2). At each visit, clinical variables were recorded, the EQ-5D was completed, and a transvaginal ultrasound was performed by the same experienced examiner.

To estimate a mean difference of 0.20 units in EQ-VAS with a SD of 0.20 and a precision of 0.10 (95% CI), 13 participants were required; we recruited 14 (p.6, l. 8–15). All variables assessed at each visit were analyzed descriptively. To analyze the changes after starting treatment, the repeat measures general linear model (GLM) was used when the quantitative variables were parametric and the Friedman test when they were nonparametric. Qualitative variables were analyzed with the Fisher or Chi-squared test to assess differences between visits. All data were entered into the study database and analyzed with the SPSS version 26.0 (IBM Corp., Armonk, NY, USA). Differences were considered statistically significant when the p-value was <0.05 . The study was approved by the Ethics Committee of Hospital de la Santa Creu i Sant Pau, Barcelona, Spain.

Statistical analysis

Analyses were performed using SPSS version 26.0 (IBM Corp., Armonk, NY, USA). To detect a mean difference of 0.20 units in EQ-VAS with an SD of 0.20 and a precision of 0.10 (95% CI), a minimum sample of 13 participants was required; 14 were included. All variables were analyzed descriptively. Changes across visits were assessed using a repeated-measures general linear model (GLM) for parametric quantitative variables and the Friedman test for nonparametric variables. Qualitative variables were analyzed with Fisher's exact or Chi-square tests. Statistical significance was set at $p < 0.05$.

Results

Of the 187 women evaluated for suspected endometriosis, 63 received combined hormonal contraceptives (CHCs) and 51 received a menopausal symptom-guided approach; the remaining women did not start treatment. NOMAC/E₂ was prescribed to the first 14 patients who accepted continuous treatment during at least 12 months.

Concomitant conditions such as uterine fibroids >3 cm, adenomyosis, inflammatory bowel disease, and irritable bowel syndrome were excluded during the initial interview. Fourteen


patients were included in the final study cohort. The mean (\pm SD) age was 36.6 ± 7.4 years (Table 1). After the first follow-up visit, 10 patients (71.4%) remained on NOMAC/E₂; after the second visit, 9 (64.3%) continued treatment (Figure 1).

Table 1. Baseline characteristics of the patients.

Baseline characteristic	n=14
Age (years)	36.6 (7.4)
History of surgery for endometriosis	3 (21.4)
Nulliparity	5 (35.7)
<ul style="list-style-type: none"> Menstrual pattern Duration of menstrual bleeding (days) Abundant menstrual bleeding Duration of dysmenorrhea (days) Intensity of dysmenorrhea (VNS) 	5 [3-10] 7 (50) 4.5 [2-10] 7.5 [3-10]
Type of endometriosis	
<ul style="list-style-type: none"> Endometrioma <ul style="list-style-type: none"> Bilateral DIE in posterior compartment <ul style="list-style-type: none"> Uterine torus Utero-sacral ligament Recto-sigma DIE in anterior compartment 	14 (100.0) 2 (14.3) 5 (35.7) 1 (7.1) 4 (28.6) 2 (14.3) 0 (0)
Body mass index (kg/m ²)	24.3 \pm 3.1
Surgical history	2 (14.3)

Data are presented as mean \pm standard deviation, median [minimum-maximum value], or frequency n (%)
 DIE, deep infiltrating pelvic endometriosis; VNS, visual numeric scale

Figure 1. Flow of participants through the study.



Reasons for treatment modification included insufficient clinical improvement in two cases in which surgery was ultimately chosen. In one patient, the switch to dienogest was motivated by persistent pain (with only slight improvement) and intermenstrual bleeding, whereas in another, mood changes prompted the transition to dienogest.

Quality-of-life outcomes demonstrated a significant improvement in both EQ-5D and EQ-VAS scores after treatment initiation (Figure 2). Significant differences were observed between baseline and both follow-up visits. Between the first and second

visits, improvement was detected only in the EQ-5D score. Clinical and ultrasound findings are summarized in Figure 3 and Table 2. Patients showed improvements in frequency and intensity of dysmenorrhea, menstrual pain, and menstrual bleeding patterns. However, no reductions were observed in the size of endometriomas or DIE lesions.

No differences in treatment response were associated with body mass index (BMI). With continuous medical therapy, 85.7% of women avoided surgery. Reported adverse effects included spotting (21.4%), decreased libido (7.1%), mood alterations (7.1%), hair loss (7.1%), and a sensation of heaviness in the legs (7.1%). Overall satisfaction with treatment was high, with median VNS scores of 4.5 at the first follow-up and 5 at the second visit.

Discussion

The results of our study suggested that treatment with a continuous regimen of NOMAC/E₂ in women presenting with dysmenorrhea and ultrasound findings suggestive of endometriosis led to a significant improvement in QoL, as measured by the EQ-5D questionnaire, after at least 6 months of uninterrupted therapy. Specifically, both the EQ-5D index score and the EQ-VAS showed statistically significant increases, indicating a marked enhancement in patients' perceived overall health status and well-being.

Endometriosis significantly impairs quality of life (QoL), as women with the condition consistently report lower QoL compared to those without the disease [5-10]. CCP, dysmenorrhea, dyspareunia, and associated physical and psychological symptoms contribute to social isolation, anxiety, depression, and reduced sexual satisfaction, collectively diminishing overall well-being [11]. Epidemiological evidence indicates that 60–70% of affected women experience physical, psychological, or social impairments that substantially affect their QoL [11-13].

Management strategies that address both symptom relief and holistic well-being are essential. Combined oral contraceptives such as NOMAC/E₂ have shown potential in alleviating pain and improving QoL, providing meaningful benefits in this population. Perceived QoL is also influenced by factors such as disease acceptance, pain severity, BMI, intimate relationships, financial burden, age, socioeconomic status, and educational level [14], emphasizing the importance of individualized treatment. Severity and duration of symptoms, as well as diagnostic delays, correlate with further reductions in QoL [7,15,13], highlighting the need for early recognition and effective management. Routine assessment using validated QoL tools such as SF-36, EHP-30, EHP-5, and EQ-5D [21,22] is crucial for evaluating treatment efficacy. Despite its relevance, research specifically analyzing QoL outcomes in endometriosis remains limited [23], underscoring the need for continued investigation into treatments that optimize both pain control and overall life quality.

In recent years, several studies confirmed that different combined oral contraceptive regimens are safe and associated with improved symptoms and reduction in endometriosis lesions [24-26].

Figure 2. QoL outcomes measured with the *EuroQoL-5D Questionnaire*.





Figure 3. Clinical outcomes.



Within broader CHC evidence, randomized and prospective data indicate that extended or continuous regimens reduce endometriosis-related pelvic pain and improve health-related QoL. Flexible/extended ethinyl-estradiol/drospirenone significantly lessens pelvic pain versus placebo and supports QoL gains in randomized trials [27]. Comparative studies of dienogest/ethinyl estradiol show that continuous regimens yield greater pain reduction and better QoL than conventional cyclic dosing [28]. For estradiol-based CHCs, the NOMAC/E₂ pill improves CCP and

QoL in observational and randomized settings, and head-to-head data show comparable pain relief to dienogest monotherapy, with some time-point advantages for dienogest on somatic QoL domains [29]. Evidence for estradiol valerate/dienogest is more limited, but available reviews describe improvements in dysmenorrhea and QoL similar to other CHCs [30].

Collectively, current literature supports CHCs—particularly in extended/continuous regimens—as effective options for

Table 2. Clinical outcomes and ultrasound findings for the patients.

Patients in follow-up	V0	V1	V2	p value*			
	n=14	n=14	n=10	Among the 3 visits	Between V0 and V1	Between V1 and V2	Between V0 and V2
Menstrual bleeding pattern	14 (100)	4 (28.6)	1 (10)	p<0.001 ^a	p<0.001 ^b	p<0.001 ^a	p=0.358 ^a
Duration (days)	5 [3-10]	0 [0-5]	0 [0-3]	p<0.001 ^c	p=0.001 ^d	p=0.005 ^d	p=0.109 ^d
Intensity				p<0.001 ^a	p<0.001 ^a	p<0.001 ^a	p=0.139 ^a
Amenorrhea	0 (0)	10 (71.4)	9 (90)				
Limited	0 (0)	3 (21.4)	1 (10)				
Normal	7 (50)	0 (0)	0 (0)				
Abundant	7 (50)	1 (7.1)	0 (0)				
Dysmenorrhea	14 (100)	4 (28.6)	0 (0)	p<0.001 ^a	p<0.001 ^b	p<0.001 ^a	p=0.049 ^a
VNS	7.5 [3-10]	0 [0-8]	0 [0-0]	p<0.001 ^c	p=0.003 ^d	p=0.005 ^d	p=0.109 ^d
Duration (days)	4.5 [2-10]	0 [0-4]	0 [0-0]	p<0.001 ^c	p=0.001 ^d	p=0.005 ^d	p=0.102 ^d
Chronic pelvic pain	3 (21.4)	5 (35.7)	3 (30.0)	p=0.75 ^a	p=0.678 ^a	p=0.665 ^a	p=1.000 ^a
VSN	5 [3-7]	6 [4-7]	4 [3-7]	p=0.638 ^c	p=1.000 ^d	p=0.317 ^d	p=0.317 ^d
Duration (days)	8 [1-20]	4 [1-21]	4 [1-14]	p=0.638 ^c	p=0.317 ^d	p=0.317 ^d	p=0.317 ^d
Days of month with pain (dysmenorrhea + CPP)	5 [2-25]	1 [0-21]	0 [0-14]	p=0.039 ^c	p=0.086 ^d	p=0.017 ^d	p=0.713 ^d
Maximum intensity of pain during month (dysmenorrhea + CPP)	7.5 [3-10]	4 [0-8]	0 [0-7]	p<0.001 ^c	p=0.011 ^d	p=0.005 ^d	p=0.141 ^d
Dyschezia (n, %)	3 (21.4)	3 (21.4)	1 (10)	p=0.764 ^a	p=0.678 ^a	p=0.665 ^a	p=1.000 ^a
VNS median [min-max]	7 [5-8]	7 [3-8]	7 [3-8]	p=0.148 ^c	p=0.414 ^d	p=0.109 ^d	p=0.180 ^d
Duration, days median [min-max]	5 [3-5]	3 [1-4]	3 [1-4]	p=0.097 ^c	p=0.180 ^d	p=0.157 ^d	p=0.180 ^d
Dyspareunia (n, %)	7 (50)	4 (28.6)	2 (20)	p=0.323 ^a	p=0.440 ^a	p=0.210 ^a	p=1.000 ^a
VNS median [min-max]	5 [2-10]	5.5 [3-9]	5.5 [3-8]	p=0.023 ^c	p=0.026 ^d	p=0.066 ^d	p=0.317 ^d
Duration, days median [min-max]	5 [4-8]	2 [2-4]	2 [2-2]	p=0.009 ^c	p=0.018 ^d	p=0.042 ^d	p=0.317 ^d
Endometrioma (n, %)	14 (100)	12 (85.7)	6 (60.0)	p=0.002 ^a	p=0.481 ^a	p=0.002 ^a	p=0.032 ^a
Maximum diameter mm (mean ± SD)	44.9 ± 21.0	41.8 ± 25.8	26.2 ± 24.8	p=0.792 ^e	p=0.278 ^f	p=0.126 ^f	p=0.187 ^f
Sum of diameters mm (mean ± SD)	59.7 ± 39.4	53.9 ± 38.6	36.7 ± 46.4	p=0.794 ^e	p=0.186 ^f	p=0.098 ^f	p=0.562 ^f
DIE (n, %)	5 (35.7)	3 (21.4)	0 (0)	p=0.009 ^a	p=0.678 ^a	p=0.005 ^a	p=0.023 ^a
Max. diameter (mm), median [min-max]	14.0 (5.4)	6.2 (9.3)	-	p=0.737 ^e	p=0.155 ^f	p=0.004 ^f	p=0.212 ^f
Spotting (n, %)	0 (0)	3 (21.4)	3 (30)	p=0.006 ^a	p=0.222 ^a	p=0.006 ^a	p=0.101 ^a
Satisfaction median [min-max]	-	4.5 [3-5]	5 [4-5]	p=1.000	-	-	-

Data are presented mean ± standard deviations, median [minimum – maximum value] or frequency n (%)

*Differences were considered statistically significant when the p value was <0.05 as determined with the ^athe Fisher exact test; ^bthe chi-square test; ^c the Friedman test; ^d the Wilcoxon test; ^e the GLM test or ^f Student's T test

CPP, chronic pelvic pain; DIE, deep infiltrating pelvic endometriosis; MBP, menstrual bleeding pattern; VNS, visual numeric scale

alleviating dysmenorrhea and pelvic pain while enhancing QoL; however, no regimen demonstrates clear, consistent superiority across outcomes, underscoring the need for individualized choices based on symptoms, tolerability, preferences, and comorbidities. Škegro et al. showed that higher pain levels correlated with poorer QoL [31]. Studies using drospirenone/EE demonstrated reductions of up to 40% in endometrioma size [32,33]. Scala et al. [34] evaluated levonorgestrel/EE in an 84/7 regimen for 12 months and observed significant reductions in symptoms and lesion size. Xholli et al. [35] reported significant reductions in endometrioma size and disappearance of lesions in nearly half of women treated with dienogest/EE for at least 12 months, along with decreases in dysmenorrhea, intermenstrual pain, and dyspareunia. Yoshino et al. [36] observed QoL improvement in Japanese women with dysmenorrhea or endometriosis-associated

pain treated with extended drospirenone/EE. QoL improvement has also been reported in women who switched from cyclic to extended regimens [37].

Regarding strengths, this prospective cohort pilot design allowed temporal evaluation of continuous NOMAC/E₂ therapy on QoL and clinical outcomes in women with ultrasound-suspected endometriosis. The validated EQ-5D instrument enabled multidimensional, patient-centered assessment, capturing changes in mobility, self-care, daily activities, pain/discomfort, and anxiety/depression. Another strength was consistency in clinical and ultrasound evaluations, all performed by a single experienced operator, minimizing inter-observer variability. The longitudinal follow-up at baseline, 6 months, and 12 months allowed assessment of both short- and intermediate-term effects.

Nonetheless, several limitations must be acknowledged. The small sample size ($n = 14$) restricts statistical power and limits generalizability. The absence of a control group prevents definitive attribution of improvements to NOMAC/E₂, as changes may reflect placebo effects or natural disease variation. As a single-center study, selection bias cannot be excluded. Although follow-up extended to 12 months, this period may be insufficient to determine long-term sustainability. Attrition was notable, with up to 40% discontinuation, which may introduce bias. Diagnosis was based on clinical and ultrasound findings rather than histological confirmation, raising the possibility of misclassification.

Conclusions

Treatment with NOMAC/E₂ in women with dysmenorrhea and ultrasound suspicion of endometriosis may significantly improve QoL as measured by EQ-5D. These findings support NOMAC/E₂ as a promising therapeutic option, although further studies are needed to confirm its benefits relative to other treatments. In our study, we did not observe reductions in lesion size, likely due to the small sample and relatively short follow-up. This highlights the importance of prioritizing patient-reported outcomes such as QoL and symptom relief, rather than relying solely on anatomical changes. Future research should include randomized controlled trials with larger cohorts and longer follow-up to confirm these results and compare different hormonal regimens.

References

- De Graaff AA, D'Hooghe TM, Dunselman GA, Dirksen CD, Hummelshoj L, WERF EndoCost Consortium; Simoens S. The significant effect of endometriosis on physical, mental and social wellbeing: results from an international cross-sectional survey. *Hum Reprod.* 2013;28(10):2677–2685.
- Moradi M, Parker M, Sneddon A, Lopez V, Ellwood D. Impact of endometriosis on women's lives: a qualitative study. *BMC Womens Health.* 2014;14:123.
- Vercellini P, Viganò P, Somigliana E, Fedele L. Endometriosis: pathogenesis and treatment. *Nat Rev Endocrinol.* 2014;10(5):261–275.
- Nnoaham KE, Hummelshoj L, Webster P, et al.; World Endometriosis Research Foundation Global Study of Women's Health consortium. Impact of endometriosis on quality of life and work productivity: a multicenter study across ten countries. *Fertil Steril.* 2011;96(2):366–373.e8.
- Gao X, Outley J, Botteman M, Spalding J, Simon JA, Pashos CL. Economic burden of endometriosis. *Fertil Steril.* 2006;86(6):1561–1572.
- Fourquet J, Báez L, Figueroa M, Iriarte RI, Flores I. Quantification of the impact of endometriosis symptoms on health-related quality of life and work productivity. *Fertil Steril.* 2011;96(1):107–112.
- Jones G, Kennedy S, Barnard A, Wong J, Jenkinson C. Development of an endometriosis quality-of-life instrument: The Endometriosis Health Profile-30. *Obstet Gynecol.* 2001;98(2):258–264.
- Jones G, Jenkinson C, Kennedy S. Evaluating the responsiveness of the Endometriosis Health Profile Questionnaire: the EHP-30. *Qual Life Res.* 2004;13(3):705–713.
- Hudelist G, Fritzer N, Thomas A, et al. Diagnostic delay for endometriosis in Austria and Germany: causes and possible consequences. *Hum Reprod.* 2012;27(12):3412–3416.
- Sepulcri Rde P, do Amaral VF. Depressive symptoms, anxiety, and quality of life in women with pelvic endometriosis. *Eur J Obstet Gynecol Reprod Biol.* 2009;142(1):53–56.
- Facchin F, Barbara G, Dridi D, et al. Mental health in women with endometriosis: searching for predictors of psychological distress. *Hum Reprod.* 2017;32(9):1855–1861.
- Markovic M, Manderson L, Warren N. Endurance and contest: women's narratives of endometriosis. *Health (London).* 2008;12(3):349–367.
- Culley L, Law C, Hudson N, et al. The social and psychological impact of endometriosis on women's lives: a critical narrative review. *Hum Reprod Update.* 2013;19(6):625–639.
- Fagervold B, Jenssen M, Hummelshoj L, Moen MH. Life after a diagnosis with endometriosis—a 15 years follow-up study. *Acta Obstet Gynecol Scand.* 2009;88(8):914–919.
- Denny E. Women's experience of endometriosis. *J Adv Nurs.* 2004;46(6):641–648.
- Soliman AM, Coyne KS, Gries KS, Castelli-Haley J, Snabes MC, Surrey ES. The effect of endometriosis symptoms on absenteeism and presenteeism in the workplace and at home. *J Manag Care Spec Pharm.* 2017;23(7):745–754.
- Ballard KD, Seaman HE, de Vries CS, Wright JT. Can symptomatology help in the diagnosis of endometriosis? Findings from a national case-control study—Part 1. *BJOG.* 2008;115(11):1382–1391.
- Spanish Fertility Society (SEF), Endometriosis Working Group. Clinical Practice Guideline: Management of Patients with Endometriosis During Reproductive Age. Spanish Fertility Society. 2018. Available at: <https://www.sefertilidad.net/index.php?seccion=biblioteca&subSeccion=guiasClinicas>.
- Vercellini P, Somigliana E, Viganò P, Abbiati A, Barbara G, Fedele L. 'Blood on the tracks' from corpora lutea to endometriomas. *BJOG.* 2009;116(4):366–371.
- Guo SW. Recurrence of endometriosis and its control. *Hum Reprod Update.* 2009;15(4):441–461.
- Simoens S, Hummelshoj L, D'Hooghe T. Endometriosis: cost estimates and methodological perspective. *Hum Reprod Update.* 2007;13(4):395–404.
- Della Corte L, Di Filippo C, Gabrielli O, et al. The burden of endometriosis on women's lifespan: a narrative overview on quality of life and psychosocial wellbeing. *Int J Environ Res Public Health.* 2020;17(13):4683.
- Zondervan KT, Becker CM, Missmer SA. Endometriosis. *N Engl J Med.* 2020;382(13):1244–1256.
- Shum LK, Bedaiwy MA, Allaire C, et al. Deep dyspareunia and sexual quality of life in women with endometriosis. *Sex Med.* 2018;6(3):224–233.
- Fernández I, Reid S, Denny E. The experience of pain and the role of psychological support in women with endometriosis: a qualitative study. *Psychol Health.* 2020;35(9):1045–1064.
- Ballweg ML. Impact of endometriosis on women's health: comparative historical data show that the earlier the onset, the more severe the disease. *Best Pract Res Clin Obstet Gynaecol.* 2004;18(2):201–218.
- Harada T, Kosaka S, Elliesen J, Yasuda M, Ito M, Momoeda M. Ethinylestradiol 20 µg/drospirenone 3 mg in a flexible extended regimen for the management of endometriosis-associated pelvic pain: a randomized controlled trial. *Fertil Steril.* 2017;108(5):798–805.
- Caruso S, Iraci M, Cianci S, Casella E, Fava V, Cianci A. Does noregestrol acetate plus 17β-estradiol oral contraceptive improve endometriosis-associated chronic pelvic pain in women? *J Womens Health (Larchmt).* 2020;29(9):1185–1191.
- Nappi RE, Serrani M, Jensen JT. Noncontraceptive benefits of the estradiol valerate/dienogest combined oral contraceptive: a review of the literature. *Int J Womens Health.* 2014;6:711–718.
- Caruso S, Cianci A, Iraci Sareri M, Panella M, Caruso G, Cianci S. Randomized study on the effectiveness of noregestrol acetate plus 17β-estradiol oral contraceptive versus dienogest oral pill in women with suspected endometriosis-associated chronic pelvic pain. *BMC Womens Health.* 2022;22(1):146.
- Škegro B, Bjedov S, Mikuš M, et al. Endometriosis, Pain and Mental Health. *Psychiatr Danub.* 2021;33(Suppl 4):632–636.
- Klein S, D'Hooghe T, Meuleman C, Dirksen C, Dunselman G, Simoens S. What is the societal burden of endometriosis-associated symptoms? a prospective Belgian study. *Reprod Biomed Online.* 2014;28(1):116–124.

33. Greaves E, Temp J, Esnal-Zufiaurre A, Mechsner S, Horne AW, Saunders PT. Estradiol is a critical mediator of macrophage-nerve cross talk in peritoneal endometriosis. *Am J Pathol.* 2015;185(9):2286–2297.
34. Scala C, Leone Roberti Maggiore U, Barra F, Venturini PL, Ferrero S. Norethindrone acetate versus extended-cycle oral contraceptive (Seasonique®) in the treatment of endometriosis symptoms: a prospective open-label comparative study. *Eur J Obstet Gynecol Reprod Biol.* 2018;222:89–94.
35. Xholli A, Filip G, Previtera F, Cagnacci A. Modification of endometrioma size during hormone therapy containing dienogest. *Gynecol Endocrinol.* 2020 Jun;36(6):545–549.
36. Yoshino O, Suzukamo Y, Yoshihara K, Takahashi N. Quality of life in Japanese patients with dysmenorrhea or endometriosis-associated pelvic pain treated with extended regimen ethinylestradiol/drospirenone in a real-world setting: a prospective observational study. *Adv Ther.* 2022;39(11):5087–5104
37. Parasar P, Ozcan P, Terry KL. Endometriosis: epidemiology, diagnosis and clinical management. *Curr Obstet Gynecol Rep.* 2017;6(1):34–41. Parasar P, Ozcan P, Terry KL. Endometriosis: epidemiology, diagnosis and clinical management. *Curr Obstet Gynecol Rep.* 2017;6(1):34–41.

Abbreviations

None.

Author contributions

Josep Perelló-Capó, Noelia Rams, Rocío Luna, Josep Estadella, Taisiia Melnychuk and Ramon Rovira conceived the study, contributed to clinical data and the main research concept.

Josep Perelló-Capó, Bruna Miarons and Natalia Teixeira: drafted the manuscript, conducted statistical analyses and performed the literature review.

Josep Perelló-Capó, Bruna Miarons, Natalia Teixeira and Rocío Luna: supported manuscript preparation, contributed additional data and participated in supervision.

Josep Perelló-Capó, Noelia Rams, Rocío Luna, Josep Estadella, Taisiia Melnychuk, Ramon Rovira: collected clinical data and reviewed the manuscript.

Josep Perelló-Capó, Noelia Rams, Josep Estadella and Taisiia Melnychuk supervised the study and provided critical revisions.

Josep Perelló-Capó, Rocío Luna and Ramon Rovira contributed with senior scientific input to the research design and the manuscript preparation, and additional statistical analysis.

All authors reviewed the manuscript and approved the final version.

Conflicts of interest

The authors declare having no conflicts of interest relevant to the content of this manuscript.

Funding

None.

Data availability statement

The data that support our findings are available from the corresponding author (Josep Perelló Capó) upon reasonable request.